



Synthesis and crystal structure of complexes of bipodal thiophosphorylated thioureas with divalent transition metals

Nail G. Zabiroy ^{a,*}, Vasiliy V. Brusko ^a, Alexander Y. Verat ^{a,1}, Dmitry B. Krivolapov ^b, Igor A. Litvinov ^b, Rafael A. Cherkasov ^a

^a Department of Chemistry, Kazan State University, Kremlevskaya Street 18, 420008, Kazan, Russia

^b Arbusov Institute of Organic and Physical Chemistry, Arbuzov St. 8, 420088, Kazan, Russia

Received 7 November 2003; accepted 1 July 2004

Available online 28 August 2004

Abstract

New complexes of bipodal thiophosphorylated thioureas $\{[(iPrO)_2P(S)NC(S)NH]_2Z\}M$, where $Z = m\text{-C}_6\text{H}_4$, $p\text{-C}_6\text{H}_4$, $M = \text{Co}$, Ni , Zn , Cd have been prepared and characterized. The crystal and molecular structures of Cd(II) , $Z = \text{CH}_2\text{CH}_2$ and Pd(II) , $Z = (\text{CH}_2\text{CH}_2)_2\text{O}$ complexes have been determined by single crystal X-ray diffraction analyses. Both complexes are binuclear in the solid state with a M_2L_2 stoichiometry. The cadmium and palladium atoms are four-coordinated and have the configuration of a slightly distorted tetrahedron for cadmium and a distorted square plane for palladium. Metal-containing macrocycles are formed as well as two spirocycles.

© 2004 Elsevier Ltd. All rights reserved.

Keywords: *P,S*-metallocycles; Supramolecular chemistry; Bipodal ligands; Crystal structures; Macrocycles; Phosphorylated urea complexes

1. Introduction

We have recently reported the syntheses and complexing properties of bipodal thiophosphorylated thioureas $[(iPrO)_2P(S)NHC(S)NH]_2Z$, where $Z = (\text{CH}_2)_2$, $(\text{CH}_2)_7$, $(\text{CH}_2)_2\text{O}(\text{CH}_2)_2$, $(\text{CH}_2)_2\text{O}(\text{CH}_2)_2\text{O}(\text{CH}_2)_2$, and some properties of the chelate complexes formed by these ligands with a series of metal ions [1,2].

In this paper, we report on the synthesis of the bis-urea $[(iPrO)_2P(S)NHC(S)NH]_2Z$ complexes with Co(II) , Ni(II) , Zn(II) and Cd(II) ($Z = m\text{-C}_6\text{H}_4$, $p\text{-C}_6\text{H}_4$), together with the X-ray study of the bis-urea $[(iPrO)_2P(S)NHC(S)NHCH_2]_2$ and the complexes with the cadmium(II) ion (**1**) ($Z = \text{CH}_2\text{CH}_2$) and the palladium(II) ion (**2**) ($Z = (\text{CH}_2\text{CH}_2)_2\text{O}$).

2. Experimental

2.1. Synthesis of $[(iPrO)_2P(S)NHC(S)NH]_2\text{-}m\text{-C}_6\text{H}_4$ (**3**)

A solution of *m*-phenylenediamine (2.163 g, 0.02 mol) in anhydrous benzene (50 ml) was added dropwise under stirring to a benzene solution (25 ml) of diisopropylthiophosphoryl isothiocyanate (9.572 g, 0.04 mol). The mixture was heated to 60 °C and stirred for 2 h. The solvent was removed in vacuum. A colourless precipitate was crystallized from benzene. Yield: 8.33 g (~71%), m.p. 122 °C. *Anal.* Calc. for $\text{C}_{20}\text{H}_{36}\text{N}_4\text{O}_4\text{P}_2\text{S}_4$: C, 40.95; H, 6.14; P, 10.58. Found: C, 41.58; H, 6.25; P, 10.30%. ¹H NMR (C_6D_6) 1.33 (d, 24H, CH_3 , ³*J*_{HH} 6.1 Hz), 4.78 (d, sept, 4H, OCH, ³*J*_{HH} 6.3 Hz, ³*J*_{POCH} 10.7 Hz), 6.86 (d, 2H, NHP(S), ²*J*_{PH} 9.8 Hz), 7.24 (t, 1H, *m*-C₆H₄, ³*J*_{HH} 8.2 Hz), 7.46 (d, d, 2H, *m*-C₆H₄, ³*J*_{HH} 7.8 Hz, ⁴*J*_{HH} 1.9 Hz), 8.14 (m, 1H, *m*-C₆H₄), 9.68 (s, 2H, C₆H₄NH). ³¹P{¹H} NMR (ppm), 53.5. IR (cm⁻¹):

* Corresponding author. Tel./fax: +7 8432 543734.

E-mail addresses: zabiroy@mi.ru (N.G. Zabiroy), averat@mail.ru (A.Y. Verat).

¹ Tel.: +7 8432 607654; fax: +7 8432 543734.